



Nov. 2010

Occupational Therapy  
Speech Therapy

**Chapel Hill Office**

205 Sage Rd, Ste 203  
Chapel Hill, NC 27514  
Phone: (919) 928-0204  
Fax: (919) 928-9423

**Durham Office**

2919 Colony Rd  
Durham, NC 27705  
Phone: (919) 489-7333  
Fax: (919) 489-0504

Serving children from  
birth to 14 years old

Using therapy to improve  
conditions such as:

- Delayed Speech
- Developmental Coordination Disorder
- Speech Impairments
- Non-communicative Speech
- Oral/Verbal Apraxia
- Language Disorders
- Feeding Difficulties
- Autism/Asperger's/PDD
- Developmental Delay
- Cerebral Palsy
- Sensory Integration Disorders
- Sensory Modulation Disorders
- Handwriting Impairments
- Fine & Gross Motor Delays

A Private, Independent  
Provider

[www.EmergeAChildsPlace.com](http://www.EmergeAChildsPlace.com)



## Social Anxiety Disorder and Overlap with Sensory Processing




Bonnie Hacker, OT  
Founder / Director

Social Anxiety Disorder is defined as a marked and persistent fear of one or more social or performance situations in which the person is exposed to unfamiliar people or to possible scrutiny by others.<sup>1</sup> It can occur in early childhood or first present in teenage years. If untreated, social anxiety disorder typically follows a chronic, unremitting course, leading to substantial challenges in career and social functioning.<sup>2,3</sup> Anxiety serves as a normal emotional response that ultimately enhances a person's ability to overcome challenges. However, for a significant percentage of children and adults, the normal anxiety response does not function properly. The distinction is simple. When anxiety proves ultimately helpful, it is normal. When anxiety is too intense or too consistent, and it ultimately interferes with needed activities, people should seek assistance.

Behaviors generated by sensory processing sensitivity overlap many behaviors of social anxiety disorder.<sup>4</sup> Research has connected sensory-processing sensitivity with anxiety, harm avoidance, and agoraphobia.<sup>5,6</sup> Sensory processing disorder (SPD) refers to the normal reception of sensory input but an exaggerated or diminished central processing of that input. Parental descriptions of a sensory processing disorder commonly include concerns over a child's level of anxiety. In children especially, fearfulness about situations compounded by the embarrassment of not fitting in sometimes results in persistent, unhealthy anxiety.

Whether SPD is a causative factor in the anxiety disorder or simply exists concurrently with anxiety disorder, moderating the SPD can result in significant improvement of the anxiety. If you know a child with anxiety problems, consider these determining questions. Does the child also demonstrate signs of SPD such as resistance to being touched, bothered by certain clothing, unpredictable temper tantrums, easily overwhelmed in groups, or too easily distracted or distressed by noises? Expert pediatric occupational therapy at Emerge - A Child's Place can help make life better for children with anxiety challenges. Our approach typically includes both "bottom up" sensory processing and "top down" cognitive tools.



**Signs of Higher than Normal Social Anxiety**

- ✿ Fear or lack of interest in new things
- ✿ Refusal to speak in social situations
- ✿ Anxiety present with peers, not just adults
- ✿ No or few friends
- ✿ Fear of separation from parents beyond normal for age
- ✿ Poor social skills

### Integrating Occupational and Speech Therapy



Among children with social anxiety, both sensory processing and speech difficulties can exacerbate the problem. For many types of challenges, Emerge provides a more comprehensive pediatric therapy approach by integrating services from both occupational therapists and speech therapists.

Please tell parents about Emerge - A Child's Place  
Believing in a Child's Potential to Flourish

## References

1. American Psychiatric Association. Diagnostic and statistical manual for mental disorders. 4. Washington, DC: APA; 1994. text revision.
2. Stein M, Kean Y. Disability and quality of life in social phobia: epidemiologic findings. *Am J Psychiatry*. 2000 Oct; 157 (10): 1606-13.
3. Stein M, Torgurd L, Walker J. Social phobia symptoms, subtypes, and severity: findings from a community survey. *Arch Gen Psychiatry*. 2000 Nov. 57 (11): 1046-52.
4. Aron E, Aron A, Davies K. Adult shyness: the interaction of temperamental sensitivity and an adverse childhood environment. 2005 Feb; 31 (2): 181-97.
5. Hofmann S, Bitran S. Sensory-processing sensitivity in social anxiety disorder: Relationship to harm avoidance and diagnostic subtypes. *Journal of Anxiety Disorders*, 2007; 21 (7): 944-54.
6. Kinnealey M, Fuiiek M. "The relationship between sensory defensiveness, anxiety, depression and perception of pain in adults." *Occupational Therapy International*; 6 (3): 195-206.